

Letters

RESEARCH LETTER

Association Between Universal Masking in a Health Care System and SARS-CoV-2 Positivity Among Health Care Workers


The coronavirus disease 2019 (COVID-19) pandemic, caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has severely affected health care workers (HCWs).¹ As a result, hospital systems began testing HCWs² and implementing infection control measures to mitigate workforce depletion and prevent disease spread.³ Mass General Brigham (MGB) is the largest health care system in Massachusetts, with 12 hospitals and more than 75 000 employees. In March 2020, MGB implemented a multipronged infection

+
Editorial

reduction strategy involving systematic testing of symptomatic HCWs and universal masking of all HCWs and patients with surgical masks.⁴ This study assessed the association of hospital masking policies with the SARS-CoV-2 infection rate among HCWs.

Methods | The institutional review board of MGB approved the study and waived informed consent. Using electronic medical records, we identified HCWs providing direct and indirect patient care who were tested for SARS-CoV-2 with reverse transcriptase-polymerase chain reaction between March 1 and April 30, 2020. The primary criterion for testing HCWs in our health care system was having symptoms consistent with SARS-CoV-2 infection. Information on the job description of each HCW was obtained by linking their record to the MGB Occupational Health Services and Human Resources databases.

Figure. Temporal Trend in Percentage Positivity of SARS-CoV-2 Testing Among HCWs



HCW indicates health care worker; MGB, Mass General Brigham; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2. All dates given are for the year 2020. The size of each data marker is proportional to the total number of SARS-CoV-2 tests performed each day over the time of the study period (x-axis), while the position of each data marker along the y-axis shows the percentage of daily test results that were positive among HCWs. The horizontal bars below the x-axis represent the timing of key interventions implemented in the state of Massachusetts and at MGB. The dotted lines represent the implementation dates of hospital policies. The study period is divided into 3 phases: a preintervention

period before implementation of universal masking of HCWs (pink), which includes March 26, the day after implementation of universal masking for HCWs, to account for HCWs who became symptomatic after business hours on March 25 and were tested on March 26; a transition period until implementation of universal masking of patients (purple) plus an additional lag period (yellow); and the intervention period (green). For the preintervention and intervention periods, daily tests were fitted by weighted nonlinear regression (curves). The change in overall slope was compared between the 2 curves to determine any statistically significant changes in trend (as shown by the P value).

We identified 3 phases during the study period: a preintervention period before implementation of universal masking of HCWs (March 1-24, 2020); a transition period until implementation of universal masking of patients (March 25–April 5, 2020) plus an additional lag period to allow for manifestations of symptoms (April 6-10, 2020), as previously defined⁵; and an intervention period (April 11-30, 2020). Positivity rates included the first positive test result for all HCWs in the numerator and HCWs who never tested positive plus those who tested positive that day in the denominator. For each HCW, any tests subsequent to their first positive test result were excluded. Using weighted nonlinear regression, we fit the best curve for the preintervention and intervention periods (based on R^2 value). The number of daily tests was used as the weight such that days with more tests had more weight in determining the curve. The overall slope of each period was calculated using linear regression to estimate the mean trend, regardless of curve shape. The change in overall slope between the preintervention and intervention periods was compared to determine any statistically significant change in mean trend, using a 2-sided $\alpha = .05$. The analysis was conducted using R version 4.0 (R Foundation).

Results | Of 9850 tested HCWs, 1271 (12.9%) had positive results for SARS-CoV-2 (median age, 39 years; 73% female; 7.4% physicians or trainees, 26.5% nurses or physician assistants, 17.8% technologists or nursing support, and 48.3% other). During the preintervention period, the SARS-CoV-2 positivity rate increased exponentially from 0% to 21.32%, with a weighted mean increase of 1.16% per day and a case doubling time of 3.6 days (95% CI, 3.0-4.5 days). During the intervention period, the positivity rate decreased linearly from 14.65% to 11.46%, with a weighted mean decline of 0.49% per day and a net slope change of 1.65% (95% CI, 1.13%-2.15%; $P < .001$) more decline per day compared with the preintervention period (Figure).

Discussion | Universal masking at MGB was associated with a significantly lower rate of SARS-CoV-2 positivity among HCWs. This association may be related to a decrease in transmission between patients and HCWs and among HCWs. The decrease in HCW infections could be confounded by other interventions inside and outside of the health care system (Figure), such as restrictions on elective procedures, social distancing measures, and increased masking in public spaces, which are limitations of this study. Despite these local and statewide measures, the case number continued to increase in Massachusetts throughout the study period,⁶ suggesting that the decrease in the SARS-CoV-2 positivity rate in MGB HCWs took place before the decrease in the general public. Randomized trials of universal masking of HCWs during a pandemic are likely not feasible. Nonetheless, these results support universal masking as part of a multipronged infection reduction strategy in health care settings.

Xiaowen Wang, MD
 Enrico G. Ferro, MD
 Guohai Zhou, PhD
 Dean Hashimoto, MD, JD
 Deepak L. Bhatt, MD, MPH

Author Affiliations: Division of Cardiovascular Medicine, Brigham and Women's Hospital, Boston, Massachusetts (Wang, Bhatt); Department of Medicine, Brigham and Women's Hospital, Boston, Massachusetts (Ferro); Center for Clinical Investigation, Brigham and Women's Hospital, Boston, Massachusetts (Zhou); Occupational Health Services, Mass General Brigham, Boston, Massachusetts (Hashimoto).

Corresponding Author: Deepak L. Bhatt, MD, MPH, Brigham and Women's Hospital, 75 Francis St, Boston, MA 02115 (dlbhatmd@post.harvard.edu).

Accepted for Publication: July 1, 2020.

Published Online: July 14, 2020. doi:10.1001/jama.2020.12897

Author Contributions: Dr Bhatt had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Drs Wang and Ferro contributed equally to this article.

Concept and design: Wang, Ferro, Hashimoto, Bhatt.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Wang, Ferro.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Wang, Zhou.

Administrative, technical, or material support: Wang, Ferro, Hashimoto.

Supervision: Hashimoto, Bhatt.

Conflict of Interest Disclosures: Dr Bhatt discloses the following relationships: advisory board: Cardax, CellProthera, Cereno Scientific, Elsevier Practice Update Cardiology, Level Ex, Medscape Cardiology, PhaseBio, PLx Pharma, Regado Biosciences; board of directors: Boston VA Research Institute, Society of Cardiovascular Patient Care, TobeSoft; chair: American Heart Association Quality Oversight Committee, NCDR-ACTION Registry Steering Committee, VA CART Research and Publications Committee; data monitoring committees: Baim Institute for Clinical Research, Cleveland Clinic, Contego Medical, Duke Clinical Research Institute, Mayo Clinic, Mount Sinai School of Medicine, Population Health Research Institute; honoraria: American College of Cardiology, Baim Institute for Clinical Research, Belvoir Publications, Duke Clinical Research Institute, HMP Global, *Journal of the American College of Cardiology*, K2P, Level Ex, Medtelligence/ReachMD, MJH Life Sciences, Population Health Research Institute, Slack Publications, Society of Cardiovascular Patient Care, WebMD; deputy editorship: *Clinical Cardiology*; research funding: Abbott, Afimmune, Amarin, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Cardax, Chiesi, CSL Behring, Eisai, Ethicon, Ferring Pharmaceuticals, Forest Laboratories, Fractyl, Idorsia, Ironwood, Ischemix, Lexicon, Lilly, Medtronic, Pfizer, PhaseBio, PLx Pharma, Regeneron, Roche, Sanofi Aventis, Synaptic, The Medicines Company; royalties: Elsevier; site coinvestigator: Biotronik, Boston Scientific, CSI, St Jude Medical, Svelte; trustee: American College of Cardiology; unfunded research: FlowCo, Merck, Novo Nordisk, Takeda. No other disclosures were reported.

Additional Contributions: We thank Stacey A. Duey, MT(ASCP), MCHP, Mass General Brigham, for assistance in extracting data from the Research Patient Data Registry, and Karen Hopcia, ScD, ANP-BC, Mass General Brigham, for assistance in extracting data from Occupational Health Services. No compensation was received for their roles.

1. Adams JG, Walls RM. Supporting the health care workforce during the COVID-19 global epidemic. *JAMA*. 2020;323(15):1439-1440. doi:10.1001/jama.2020.3972

2. Hunter E, Price DA, Murphy E, et al. First experience of COVID-19 screening of health-care workers in England. *Lancet*. 2020;395(10234):e77-e78. doi:10.1016/S0140-6736(20)30970-3

3. Black JRM, Bailey C, Przewrocka J, Dijkstra KK, Swanton C. COVID-19: the case for health-care worker screening to prevent hospital transmission. *Lancet*. 2020;395(10234):1418-1420. doi:10.1016/S0140-6736(20)30917-X

4. Klompas M, Morris CA, Sinclair J, Pearson M, Shenoy ES. Universal masking in hospitals in the Covid-19 era. *N Engl J Med*. 2020;382(21):e63. doi:10.1056/NEJMp2006372

5. Sen S, Karaca-Mandic P, Georgiou A. Association of stay-at-home orders with COVID-19 hospitalizations in 4 states. *JAMA*. 2020;323(24):2522-2524. doi:10.1001/jama.2020.9176

6. Massachusetts Department of Public Health COVID-19 dashboard—April 30, 2020. Accessed June 27, 2020. <https://www.mass.gov/doc/covid-19-dashboard-april-30-2020/download>